

DCD Early Identification and Intervention Sheet



Advocacy Toolkit

What is Developmental Coordination Disorder (DCD)?

Developmental Coordination Disorder (DCD) is a chronic motor skill disorder seen in children and youth, which significantly affects activities of daily living, school performance, and leisure activities.^{1,2} In order to meet DCD diagnostic criteria, the motor deficits must not be the result of any other known neurological or medical condition (such as cerebral palsy or a neurodegenerative disorder).¹

The disorder is diagnosed using DSM-5 criteria:³

- A.** Acquisition and execution of coordinated motor skills are substantially below what would be expected given the child's age and opportunity for skill learning and use. Difficulties may be seen as clumsiness, inaccuracy, or slowness of performance of motor skills (e.g., catching a ball, using scissors, printing or handwriting, riding a bicycle, or participating in sports).
- B.** The motor skills deficit significantly and persistently interferes with activities of daily living and impacts school productivity, vocational skills, leisure activities, and play.
- C.** The onset of symptoms is in the early developmental period.
- D.** The motor skills deficit is not better explained by intellectual disability, visual impairment, or a neurological or medical condition affecting movement.

Can DCD be identified before the age of 5 years?

DCD is usually evident early on in a child's life but is not typically diagnosed before age 5.¹ However, preschoolers (aged 3 to 5 years) who show significant motor impairments (despite having had ample opportunities for learning and with other causes of motor delay ruled out) can receive a DCD diagnosis based on the findings from at least two longitudinal assessments (e.g., repeated administration of the MABC-2 at least 3 months apart).¹

Why should we focus on early identification of DCD?

Parents who seek help for young children often experience greater delays and difficulties getting a DCD diagnosis, as compared to older children, which may contribute to higher stress levels.³ Earlier diagnoses and interventions for children at risk of DCD may change the developmental trajectory for these children and positively contribute to family functioning and overall well-being.

What are the risk factors for and early signs of DCD?

While the diagnosis of DCD may not be appropriate in the early years, occupational therapists have an important role in identifying children who may be at risk of DCD. Key risk factors for DCD include prematurity,⁴ low birth weight,⁴ male sex,⁴ autism,⁵ and significant speech and/or language difficulties that persist at 4-5 years of age.⁶

Several early signs for possible DCD have been proposed, largely based on clinical impressions, history taking, and parent report.^{3,6-9} Commonly identified signs include poor performance in activities of daily living (ADLs) that require motor coordination (e.g., dressing, utensil use), poor balance, clumsiness, poor motor sequencing, spatial awareness problems (e.g., poor figure ground skills and falling/bumping into others), poor visual tracking, immature grasp, poorly established hand dominance, and slower, less accurate movements. Motor milestones may or may not be delayed, but there is a history of difficulty learning age-appropriate motor skills.

Who can help to identify children with/at risk of DCD before the age of 5 years?

Parents play a critical role in the initial identification of motor difficulties for their children. Half of parents surveyed identified concerns prior to their child's 3rd birthday and sought help approximately 1.5 years after they first started to have concerns.³ A smaller study found that nearly all the parents in an interview sample had identified concerns by age 4.¹⁰ Early parent anxieties regarding poor motor development and behaviour are often confirmed by **early educators** in daycare or preschool settings.^{11,12} These early educators are well poised for naturalistic observations of children in play and can be guided in identifying those most at risk of motor coordination difficulties;⁷ most notably by **occupational therapists**¹³ and **physical therapists**.¹⁴ More attention has also been given to the potential role of **speech language pathologists** as early identifiers of children at risk for DCD given the high co-occurrence of speech and language impairments and motor impairments.⁶ **Physicians**¹⁵ are positioned to potentially identify children through well-baby and annual check-ups, but they will likely need a formal assessment of motor skills (Criterion A) and documentation of the impact on the motor skills deficit on daily life (Criterion B) (See Letter to the Doctor template [<http://bit.ly/2D8IDEY>]).

What assessment tools can be used for early identification of children with/at risk of DCD?

Criterion A: The most commonly used measure is the **Movement Assessment Battery for Children-2** (MABC-2).¹⁶ The MABC-2 has been shown to have excellent sensitivity (identifying children at risk of DCD) but also many false positives at age 3 years.¹⁷ A cut-off score of $\leq$ 5th percentile is used for children $\leq$ 5 years of age, usually over 2 assessments at least 3 months apart.¹ Children at risk of DCD may be identified in infancy. For example, a retrospective study of preterm children diagnosed with DCD at 4.5 years found that these children scored more poorly on ALL early motor assessments, from as young as 4 months of age, compared to children who did not have motor problems.¹⁸ These assessments include the **Movement Assessment of Infants**,¹⁹ the **Bayley Test of Infant and Toddler Development – 3rd ed**^{3,20} and the **Peabody Developmental Motor Scale – 2nd ed**.²¹ The **Alberta Infant Motor Scale**²² has also been used to identify preterm infants at risk of later motor impairments.^{23,24}

Criterion B: The most studied and the only questionnaire validated in Canada is the **Little Developmental Coordination Disorder Questionnaire (Little DCDQ)** (available for \$50 CAD at <http://www.dcdq.ca/little-dcdq-ca.html>).²⁵ The Little DCDQ is a parent-report questionnaire designed specifically to identify children (3 years to 4 years 11 months) at risk of DCD. Other questionnaires include the DCDQ²⁶

(<http://www.dcdq.ca>, for children aged 5 years+), and the **Early Years Movement Skills Checklist** (a parent- and teacher-report measure)²⁷ developed in the United Kingdom.

What treatment approaches can be used for children with/at risk of DCD?

Individual intervention is generally recommended for younger children,¹ but evidence is emerging for group-level interventions.²⁸⁻³¹ The following treatment approaches show promise (mostly based on Level V best evidence, e.g., case reports, single subject research designs, pre/post designs with no control group) for younger children with/at risk of DCD:

- Modified Cognitive Orientation to Occupational Performance (CO-OP)³²⁻³⁴
- Task-specific intervention and parent education³¹
- Occupational Performance Coaching (teacher) and play-based intervention (child)³⁵
- Motor Magic (occupational therapy embedded in the curriculum)^{29,30}
- Fundamental Movement Skills²⁸
- Animal Fun³⁶ (specific results for children with DCD have not yet been published)
- Strength training³⁷

Where can I learn more?

- Results of scoping review of early identification and early intervention of children with/at risk of DCD will soon be submitted for publication (Lee & Zwicker, in preparation)
- Evidence on DCD assessment/diagnosis and management [Evidence for Practice (E4P) Synthesis]: <http://www.childdevelopment.ca/DCDAdvocacyToolkit/DCDAdvocacyToolkitEvidenceforPracticeSummary.aspx>
- Advocating for a DCD diagnosis (information sheet): <http://www.childdevelopment.ca/DCDAdvocacyToolkit/DCDAdvocacyToolkitResources.aspx>
- Review of standardized motor assessments: <http://www.therapybc.ca/eLibrary/resources.php>

This document was prepared in March 2018 and will be updated as new evidence emerges.

References

1. Blank R, Smits-Engelsman B, Polatajko H, Wilson P. European Academy for Childhood Disability (EACD): Recommendations on the definition, diagnosis, and intervention of developmental coordination disorder (long version). *Dev Med Child Neurol*. 2012;54:54-93.
2. Chung A, Callanen A. Clinical Review: Developmental Coordination Disorder (Occupational Therapy). CINAHL Information Systems, Rehabilitation Reference Center, 2016.
3. Alonso Soriano C, Hill EL, Crane L. Surveying parental experiences of receiving a diagnosis of developmental coordination disorder (DCD). *Res Dev Disabil*. 2015;43-44:11-20.
4. Zwicker JG, Yoon SW, Mackay M, et al. Perinatal and neonatal predictors of developmental coordination disorder. *Arch Dis Children – Fetal and Neonatal Ed*, 2013;98:118-122.
5. Provost B, Lopez BR, Heimerl S. A comparison of motor delays in young children: Autism spectrum disorder, developmental delay, and developmental concerns. *J Autism Dev Dis*. 2006;37:321-328.
6. Gaines R, Missiuna C. Early identification: Are speech/language-impaired toddlers at increased risk for developmental coordination disorder?. *Child Care Health Dev*, 2007;33:325-332.
7. Cailloux F, Spica O. L'enfant dyspraxique en maternelle: Les signes d'appel, présentation d'une étude exploratoire concernant un outil de repérage, quelques pistes d'adaptations pédagogiques. = dyspraxie children in the école maternelle (3-6 year olds): Presentation of an exploratory study into a screening tool-- some avenues for pedagogic adaptations. A.N.A.E. Approche Neuropsychologique Des Apprentissages Chez l'Enfant, 2006;18:172-178.
8. Hertz J, Estes B. Developmental dyspraxia and developmental coordination disorder. In A. S. Davis, & A. S. Davis (Eds.) *Handbook of Pediatric Neuropsychology* (pp. 593-602). New York, NY, US: Springer Publishing Co., 2011.
9. Kirby A. (2001). Co-ordination problems. *Community Practitioner*, 2001;74:294-296.
10. Missiuna C, Moll S, King S, King G, Law M. A trajectory of troubles: Parents' impressions of the impact of developmental coordination disorder. *Phys Occup Ther Pediatr*. 2007;27: 81-101.
11. Portwood M. Developmental dyspraxia: Identification and intervention. A manual for parents and professionals (2nd ed.). London: David Fulton Publishers, 1999.
12. Boon M. Helping children with dyspraxia. London: Jessica Kingsley Publishers, 2001.

13. Missiuna C, Pollock N, Campbell W, et al. Using an innovative model of service delivery to identify children who are struggling in school. *Br J Occup Ther*. 2017;80:145-154.
14. Pridham EJ. Identification and management of four-year-old children at risk for developmental coordination disorder (DCD). Unpublished doctoral dissertation. University of South Australia, Adelaide, South Australia, 2013.
15. Harris SR, Mickelson ECR, Zwicker JG. Diagnosis and management of developmental coordination disorder. *Can Med Assoc J*. 2015;187:659-665.
16. Henderson SE, Sugden D A, Barnett AL. Movement Assessment Battery for Children-2nd ed. [MABC-2]. London: Psychological Corporation, 2007.
17. Kwok C, Mackay M, Agnew JA, Synnes A, Zwicker JG. Can the Movement Assessment Battery for Children-2 reliably identify developmental coordination disorder in 3-year-old preterm children? *Res Dev Disabil*. In press
18. Isberg D, Hare S, Mackay M, Synnes A, Grunau RE, Miller SP, Zwicker JG. Can developmental coordination disorder be predicted by early motor performance? Canadian Association of Occupational Therapists Conference, May 27-30, 2015, Winnipeg, MB.
19. Chandler LS, Andrews MS, Swanson MW. Movement Assessment of Infants: A Manual. Larson AH (ed). Seattle, WA: Child Development and Mental Retardation Center, University of Washington, 1980.
20. Bayley N. Bayley Scales of Infant and Toddler Development (3rd ed.). San Antonio, TX: Psychological Corporation, 2006.
21. Folio MR, Fewell RR. Peabody Developmental Motor Scales Examiner's Manual, 2nd ed. Austin, TX: Pro-Ed, 2000.
22. Piper MC, Pinnell LE, Darrach J, et al. Construction and validation of the Alberta Infant Motor Scale (AIMS). *Can J Public Health*, 1992;83, S46-50.
23. Spittle A, Loreface LE, Anderson PJ, et al. The validity of the Alberta Infant Motor Scale in predicting motor performance at 4.5 years old in very preterm infants. *Physiother*. 2011;97:eS1166-eS1167.
24. Spittle AJ, Anderson PJ, Lee KJ, et al. Motor assessments in the first year of life for preterm infants predict motor outcomes at preschool age. *Dev Med Child Neurol*. 2014;56:95-96.
25. Wilson BN, Creighton D, Crawford SG, et al. Psychometric properties of the Canadian Little Developmental Coordination Disorder Questionnaire for preschool children. *Phys Occup Ther Pediatr*. 2015;35:116-131.
26. Wilson BN, Crawford SG, Green D, Roberts G, Aylott A, & Kaplan B. Psychometric properties of the revised Developmental Coordination Disorder Questionnaire. *Phys Occup Ther Pediatr*. 2009;29(2):182-202.
27. Chambers ME, Sugden DA. Early years movement skills – Description, diagnosis and intervention. London: Wiley, 2006.
28. Bardid F, Deconinck FJA, Descamps S, et al. (2013). The effectiveness of a fundamental motor skill intervention in pre-schoolers with motor problems depends on gender but not environmental context. *Res Dev Disabil*. 2013;34:4571-4581.
29. Priest N. 'Motor magic': Evaluation of a community capacity-building approach to supporting the development of preschool children. *Aus Occup Ther J*. 2006;53:220-232.
30. Priest N, Waters E. 'Motor magic': Evaluation of a community capacity-building approach to supporting the development of preschool children (part 2). *Aus Occup Ther J*. 2007;54:140-148.
31. Kane KJ, Staples KL. A group motor skills program for children with coordination difficulties: Effect on fundamental movement skills and physical activity participation. *Phys Occup Ther Pediatr*. 2016;36:28-45.
32. Taylor S, Fayed N, Mandich A. CO-OP intervention for young children with developmental coordination disorder. *OTJR: Occup Participation Health*. 2007;27(4):124-130.
33. Liu CH. The effects of Cognitive Orientation to daily Occupational Performance approach for motor coordination disorder. 94th Annual Conference of the American Congress of Rehabilitation Medicine, ACRM 2017, Arch Phys Med Rehab. 2017;98(10):e117.
34. Zwicker JG, Mackay M, Cuthbert D, Synnes A. Effectiveness of Cognitive Orientation to Occupational Performance (CO-OP) in very preterm preschool-age children with developmental coordination disorder: A pilot study. Occupational Science & Occupational Therapy Research Initiative Fund Annual Report [unpublished], 2017.
35. Kennedy-Behr A, Rodger S, Graham F, Mickan, S. Creating enabling environments at preschool for children with developmental coordination disorder. *J Occup Ther, Schools Early Int*. 2013;6:301-313.
36. Piek JP, McLaren S., Kane R, et al. Does the Animal Fun program improve motor performance in children aged 4-6 years? *Hum Mov Sci*. 2013;32:1086-1096.
37. Kaufman LB, Schilling DL. Implementation of a strength training program for a 5-year-old child with poor body awareness and developmental coordination disorder. *Phys Ther*. 2007;87:455-467.



[Access the full Toolkit here](#)



This document was developed by Giovanna Boniface, Stephanie Glegg, Ivonne Montgomery, and Dr. Jill Zwicker with funding support from a Michael Smith Foundation for Health Research (MSFHR) REACH Award and Sunny Hill Health Centre for Children. Stephanie Glegg is supported by a CIHR Vanier Canada Graduate Scholarship, the Canadian Child Health Clinician Scientist Program, a UBC Public Scholars Award and Four-Year Fellowship, and the Sunny Hill Foundation. Dr. Zwicker is a MSFHR Scholar and is funded by the Canadian Child Health Clinician Scientist Program, BC Children's Hospital Research Institute, Sunny Hill Foundation, and Canadian Institutes of Health Research.